

May 14 to May 20, 2017 (Week 20)


Overall Summary

- Overall, influenza activity continues to decline slowly in Canada.
- In week 20, influenza B accounted for the majority of influenza activity in Canada, with 75% or more of reported influenza laboratory detections, hospitalizations and outbreaks due to influenza B.
- This increase in influenza B activity is expected as influenza B often appears later in the flu season.
- To date, the majority of laboratory detections, hospitalizations and deaths have been among adults aged 65+ years.
- The current FluWatch report is the last weekly report for the 2016-2017 season. During the summer, monthly reports will be published starting on June 23, 2017. We will continue to monitor influenza and other respiratory infections via the [RVDSS report](#), published every Thursday.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

In week 20, 19 regions across eight provinces and territories reported no influenza or influenza-like illness activity. Sporadic influenza activity was reported in 25 regions across 11 provinces and territories. Localized activity was reported in nine regions across five provinces. For more details on a specific region, click on the map.

Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 20




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in the mapping feature found in the [Weekly Influenza Reports](#).

Laboratory Confirmed Influenza Detections

In week 20, the number (245) and the percentage (6.8%) of tests positive for influenza decreased from the previous week. Influenza B was the most common type of influenza detected in all jurisdictions in Canada. Since week 16, the percentage of tests positive for influenza B has plateaued (6.0% to 6.9%) and has not yet begun a clear decline. Overall in week 20, influenza B accounted for 86% of total detections. Influenza B detections are within expected levels compared to the same time period in recent seasons. For data on other respiratory virus detections, see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada (PHAC) website.


Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2016-17, Week 20



The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

To date this season, 38,570 laboratory confirmed influenza detections have been reported, of which 90% have been influenza A. Influenza A(H3N2) has been the most common subtype detected this season. For more detailed weekly and cumulative influenza data, see the text descriptions for Figures 2 and 3 or the [Respiratory Virus Detections in Canada Report](#).

Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province/territory, Canada, 2016-17, Week 20



To date this season, detailed information on age and type/subtype has been received for 26,534 laboratory-confirmed influenza cases (Table 1). Among cases with reported age and type/subtype information, adults aged 65+ accounted for half of the reported influenza cases. Adults aged 65+ have predominantly been affected by influenza A accounting for 51% of influenza A detections. Influenza B, while much smaller in number, is mainly affecting individuals less than 65 years of age.

Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting¹, Canada, 2016-17, Week 20

Age groups (years)	Week (May 14, 2017 to May 20, 2017)					Cumulative (August 28, 2016 to May 20, 2017)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) ³		A Total	A(H1) pdm09	A(H3)	A (UnS) ³		Total	#
0-4	5	0	<5	<5	<5	2235	15	830	1390	246	2481	9%
5-19	0	0	0	0	15	2196	11	1077	1108	473	2669	10%
20-44	<5	0	0	<5	13	3406	23	1801	1582	480	3886	15%
45-64	<5	0	0	<5	24	3893	21	1954	1918	645	4538	17%
65+	>9	0	9	<5	42	12060	10	5441	6609	900	12960	49%
Total	>21	0	>9	12	>94	23790	80	11103	12607	2744	26534	100%
Percentage²	18%	0%	x%	x%	82%	90%	0%	47%	53%	10%		

¹Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. Cumulative data include updates to previous weeks.

²Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

³UnS: unsubtyped: The specimen was typed as influenza A, but no result for subtyping was available.

x - Supressed to prevent residual disclosure

Specimens from NT, YT, and NU are sent to reference laboratories in the provinces


Syndromic/Influenza-like Illness Surveillance

Healthcare Professionals Sentinel Syndromic Surveillance

In week 20, 0.9% of visits to healthcare professionals were due to influenza-like illness.

Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, 2016-17

Number of Sentinels Reporting Week 20: 94



Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data are compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.


Are you a primary healthcare practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel? Please visit our [Influenza Sentinel page](#) for more details.

Influenza Outbreak Surveillance

In week 20, nine laboratory-confirmed influenza outbreaks were reported. Of the four outbreaks with known strains or subtypes, three were due to influenza B and one was due to influenza A(H3N2). An additional outbreak due to ILI was reported in a school.

To date this season, 1,176 outbreaks have been reported and the majority (66%) have occurred in LTC facilities. A total of 82 outbreaks (7%) due to influenza B have been reported. Compared to the same period in the most recent previous A(H3N2) predominant season (2014-15), 1,724 outbreaks were reported, of which 74% occurred in LTC facilities.

Figure 5 – Number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2016-17, Week 20



¹All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of this report.

Provincial/Territorial Influenza Hospitalizations and Deaths

In week 20, 46 influenza-associated hospitalizations were reported by participating provinces and territories*, a slight decrease from 52 hospitalizations reported in the previous week. In week 20, 80% of reported hospitalizations were due to influenza B and 61% occurred in adults 65+. Additionally, less than five intensive care unit (ICU) admissions and five deaths were reported.

To date this season, 6,335 hospitalizations have been reported, of which 89% were due to influenza A. Among cases for which the subtype of influenza A was reported, 99% were influenza A(H3N2). Adults 65+ accounted for 88% of the hospitalizations. A total of 259 ICU admissions and 371 deaths have been reported. The majority of deaths (88%) were reported in adults aged 65+ years.

Table 2 – Cumulative number of hospitalizations, ICU admissions and deaths by age and influenza type reported by participating provinces and territories, Canada, 2016-17, Week 20

Age Groups (years)	Cumulative (August 28, 2016 to May 20, 2017)						
	Hospitalizations			ICU Admissions		Deaths	
	Influenza A Total	Influenza B Total	Total [# (%)]	Influenza A and B Total	%	Influenza A and B Total	%
0-4	444	80	524 (8%)	18	7%	<5	x%
5-19	240	83	323 (5%)	20	8%	<5	x%
20-44	294	43	337 (5%)	24	9%	5	1%
45-64	761	118	879 (14%)	78	30%	35	9%
65+	3917	355	4272 (68%)	119	46%	326	88%
Total	5656	679	6335 (100%)	259	100%	371	100%

x: Suppressed to prevent residual disclosure

*Note: Influenza-associated hospitalizations are not reported to PHAC by BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not distinguished among hospital admissions reported from ON. The hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

Sentinel Hospital Influenza Surveillance

Pediatric Influenza Hospitalizations and Deaths

In week 20, 13 laboratory-confirmed influenza-associated pediatric (≤ 16 years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network. All hospitalizations were associated with influenza B and 69% were in children over the age of 5 years.

To date this season, 569 laboratory-confirmed influenza-associated pediatric hospitalizations were reported by the IMPACT network. Children aged 0-23 months accounted for approximately 37% of hospitalizations and influenza A accounted for 79% of the reported hospitalizations. Among the 117 hospitalizations due to influenza B, 64 (54%) were in children over the age of 5 years. In comparison, children over the age of 5 years accounted for 33% of influenza A hospitalizations. Additionally, 93 intensive care unit (ICU) admissions have been reported. A total of 64 ICU cases (69%) reported at least one underlying condition or comorbidity. Less than five deaths have been reported this season.

Figure 6 – Cumulative numbers of pediatric hospitalizations (≤ 16 years of age) with influenza by age-group reported by the IMPACT network, Canada, 2016-17, Week 20





Figure 7 – Number of pediatric hospitalizations (≤ 16 years of age) with influenza reported by the IMPACT network, by week, Canada, 2016-17, Week 20



The shaded area represents the maximum and minimum number of cases reported by week from seasons 2010-11 to 2015-16

The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated pediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Adult Influenza Hospitalizations and Deaths

Surveillance for the 2016-2017 influenza season ended on April 30th, 2017.

This season, 1,535 laboratory-confirmed influenza-associated adult (≥ 20 years of age) hospitalizations have been reported by CIRN. Influenza A accounted for 92% of hospitalizations. Adults aged 65+ accounted for 78% of hospitalizations. A total of 143 intensive care unit (ICU) admissions have been reported. Among ICU cases with available data, 126 cases (88%) reported at least one underlying condition or comorbidity. The median age of patients admitted to the ICU was 71 years. Approximately 86 deaths have been reported this season, the majority in adults aged 65+. The median age of reported deaths was 85 years.

Figure 8 - Cumulative numbers of adult hospitalizations (≥ 20 years of age) with influenza by type and age-group reported by CIRN, Canada, 2016-17, Week 20





Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza by age-group (≥ 20 years of age) reported by CIRN, Canada 2016-17, Week 20



The number of hospitalizations reported through CIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Influenza Strain Characterizations

During the 2016-17 influenza season, the National Microbiology Laboratory (NML) has characterized 2,050 influenza viruses [1565 A(H3N2), 48 A(H1N1), 437 influenza B]. All but one influenza A virus (n=1613) and 95 influenza B viruses characterized were antigenically or genetically similar to the vaccine strains included in both the trivalent and quadrivalent vaccines. Three hundred and forty-two influenza B viruses were similar to the strain which is only included in the quadrivalent vaccine.

Table 3 – Influenza strain characterizations, Canada, 2016-17, Week 20

Strain Characterization Results ¹	Count	Description
Influenza A (H3N2)		
Antigenically A/Hong Kong/4801/2014-like	369	Viruses antigenically similar to A/Hong Kong/4801/2014, the A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.
Genetically ² A/Hong Kong/4801/2014-like	1195	Viruses belonging to genetic group 3C.2a. A/Hong Kong/4801/2014-like virus belongs to genetic group 3C.2a and is the influenza A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine. Additionally, genetic characterization of the 369 influenza A (H3N2) viruses that underwent HI testing determined that 306 viruses belonged to genetic group 3C.2a and 63 viruses belonged to genetic group 3C.3a. The majority of viruses belonging to genetic group 3C.3a are inhibited by antisera raised against A/Hong Kong/4801/2014 ³ .
Antigenically A/Indiana/10/2011-like ⁴	1	Viruses antigenically similar to A/Indiana/10/2011, a candidate H3N2v vaccine virus.
Influenza A (H1N1)		
A/California/7/2009-like	48	Viruses antigenically similar to A/California/7/2009, the A(H1N1) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
Influenza B		
B/Brisbane/60/2008-like (Victoria lineage)	95	Viruses antigenically similar to B/Brisbane/60/2008, the influenza B component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
B/Phuket/3073/2013-like (Yamagata lineage)	342	Viruses antigenically similar to B/Phuket/3073/2013, the additional influenza B component of the 2016-17 Northern Hemisphere quadrivalent influenza vaccine.

¹The NML receives a proportion of the influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Strain characterization data reflect the results of hemagglutination inhibition (HI) testing compared to the reference influenza strains recommended by [WHO](#).

²Determined by sequence analysis

³[WHO](#) - Recommended composition of the influenza virus vaccines for use in the 2016-17 northern hemisphere influenza season.

⁴Detected in epidemiological week 50. For more details, see [Week 50 report](#)

Antiviral Resistance

During the 2016-17 season, the National Microbiology Laboratory (NML) has tested 1,122 influenza viruses for resistance to oseltamivir, 1,124 influenza viruses for resistance to zanamivir and 247 influenza viruses for resistance to amantadine. All but two influenza A(H3N2) viruses were sensitive to oseltamivir and all viruses were sensitive to zanamivir. All 228 influenza A viruses were resistant to amantadine (Table 4).

Table 4 – Antiviral resistance by influenza virus type and subtype, Canada, 2016-17, Week 20

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	748	2 (0.3%)	747	0 (0%)	206	206 (100%)
A (H3N2v)	1	0 (0%)	1	0 (0%)	1	1 (100%)
A (H1N1)	42	0 (0%)	41	0 (0%)	40	40 (100%)
B	331	0 (0%)	335	0 (0%)	NA ¹	NA ¹
TOTAL	1122	2 (0.2%)	1124	0 (0%)	247	247 (100%)

¹NA: Not Applicable

Provincial and International Influenza Reports

- [World Health Organization influenza update](#)
- [World Health Organization FluNet](#)
- [WHO Influenza at the human-animal interface](#)
- [Centers for Disease Control and Prevention seasonal influenza report](#)
- [European Centre for Disease Prevention and Control - epidemiological data](#)
- [South Africa Influenza surveillance report](#)
- [New Zealand Public Health Surveillance](#)
- [Australia Influenza Report](#)
- [Pan-American Health Organization Influenza Situation Report](#)
- [Alberta Health – Influenza Surveillance Report](#)
- [BC - Centre for Disease Control \(BCCDC\) - Influenza Surveillance](#)
- [New Brunswick – Influenza Surveillance Reports](#)
- [Newfoundland and Labrador – Surveillance and Disease Reports](#)
- [Nova Scotia - Flu Information](#)
- [Public Health Ontario – Ontario Respiratory Pathogen Bulletin](#)
- [Manitoba – Epidemiology and Surveillance – Influenza Reports](#)
- [Saskatchewan – influenza Reports](#)
- [PEI – Influenza Summary](#)

FluWatch Definitions for the 2016-2017 Season

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

Influenza-like-illness (ILI): Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI/Influenza outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.

Influenza/ILI Activity Levels

1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

3 = Localized: (1) evidence of increased ILI* ;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

4 = Widespread: (1) evidence of increased ILI*;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

** More than just sporadic as determined by the provincial/territorial epidemiologist.*

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This [report](#) is available on the Government of Canada Influenza webpage. Ce rapport est disponible dans les deux langues officielles.