

# RESPIRATORY VIRUS DETECTION REPORT

April 3 to April 9, 2022  
(Surveillance Week 2022-14)



In this Respiratory Virus Report, the number of detections of coronavirus reflects only seasonal human coronaviruses, not the novel pandemic coronavirus (SARS-CoV-2) that causes COVID-19. For information on COVID-19, see the [Coronavirus disease \(COVID-19\): Outbreak update](#).

## Overall Summary

Detections of seasonal respiratory viruses have increased since February 2022. Activity of all respiratory viruses is within or below expected levels for this time of year.

- Respiratory syncytial virus (414 detections; 4.3% positive) is the most common seasonal respiratory virus detected in Canada, but activity is stable and below expected levels.
- A total of 363 influenza detections were reported (362 influenza A, 1 influenza B). Activity of influenza A has increased in recent weeks, and is approaching expected levels for this time of year.
- Detections of other respiratory viruses are low.
- The number of tests performed for all viruses were within or above expected levels.
- 32/34 sentinel laboratories reported surveillance data.

## Abbreviations

Flu: Influenza

Flu A(H1N1)pdm09: Influenza A(H1N1)pdm09

Flu A(H3): Influenza A(H3N2)

Flu A (UnS): Influenza A(Unsubtyped)

RSV: Respiratory syncytial virus

HPIV: Human parainfluenza virus

HMPV: Human metapneumovirus

ADV: Adenovirus

EV/RV: Enterovirus/Rhinovirus

HCoV: Human coronavirus

Atl: Atlantic Canada

BC: British Columbia

Can: Canada

EORLA: Eastern Ontario Regional Laboratory Association

HSC: Health Sciences Centre

N.C.: Data not collected

N.R.: Data not reported for current week

ON: Ontario

PR: Prairies

P.H.O.L.: Public Health Ontario Laboratory


QC: Quebec

Terr: Territories of Canada

UHN: University Health Network



**Figure 1: Positive respiratory virus tests (%) reported by participating laboratories in Canada by surveillance week compared to average and range from 2014-2015 to 2019-2020 season**



Shaded area represents the maximum and minimum % of respiratory virus tests positive reported by week from seasons 2014-2015 to 2019-2020

Note: Data from week 11 of the 2019-2020 season onwards are excluded from the historical comparison due to the COVID-19 pandemic

Note: % Tests Positive scale changes across the dashed line

<sup>a</sup> Human coronavirus excludes human coronaviruses SARS-CoV, MERS-CoV and SARS-CoV-2; Includes seasonal human coronaviruses HCoV-229E, HCoV-OC43, HCoV-NL63, HCoV-HKU1.





**Figure 2: Number of positive respiratory virus tests reported by participating laboratories in Canada by surveillance week**




Figure 3: Positive Influenza (Flu) tests (%) in Canada by region by surveillance week




Figure 4: Positive Respiratory syncytial virus (RSV) tests (%) in Canada by region by surveillance week




Figure 5: Positive Human parainfluenza virus (HPIV) tests (%) in Canada by region by surveillance week






Figure 6: Positive Adenovirus (ADV) tests (%) in Canada by region by surveillance week




Figure 7: Positive Human metapneumovirus (HMPV) tests (%) in Canada by region by surveillance week




Figure 8: Positive Enterovirus/Rhinovirus (EV/RV) tests (%) in Canada by region by surveillance week




Figure 9: Positive Human coronavirus (HCoV) tests (%) in Canada by region by surveillance week

